UNITED STATES DEPARTMENT OF THE INTERIOR, Fred A. Seaton, *Secretary*
FISH AND WILDLIFE SERVICE, Arnie J. Suomela, *Commissioner*

TREATMENT OF SULFONAMIDE-RESISTANT FURUNCULOSIS IN TROUT AND DETERMINATION OF DRUG SENSITIVITY

By S. F. Snieszko and G. L. Bullock

FISHERY BULLETIN 125
From Fishery Bulletin of the Fish and Wildlife Service
VOLUME 57

UNITED STATES GOVERNMENT PRINTING OFFICE • WASHINGTON : 1957

For sale by the Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C.
Price 15 cents

ABSTRACT

Treatment of furunculosis with sulfonamides was unsuccessful when the outbreak was caused by sulfonamide-resistant variants of the pathogen. The pathogen was sensitive to chloramphenicol and treatment with this antibiotic resulted in greatly reduced mortalities. Losses were also greatly affected by the strain-specific resistance or susceptibility of the experimental brook trout to furunculosis.

Sensitivity or resistance of *Aeromonas salmonicida* to sulfonamides and antibiotics was determined in vitro, using various liquid and solid bacteriological media. A simple method for the determination in vitro of sulfonamide- and antibiotic-sensitivity of *A. salmonicida* is described for field use. A revised method of treatment of furunculosis is also outlined.

CONTENTS

	Page
Introduction	555
Materials and methods	555
Results	556
Treatment with sulfonamides and chloramphenicol	556
Sensitivity tests with antibiotics	558
Sensitivity tests with sulfonamides	558
Discussion	562
Recommendations	563
Summary	563
Literature cited	564

TREATMENT OF SULFONAMIDE-RESISTANT FURUNCULOSIS IN TROUT AND DETERMINATION OF DRUG SENSITIVITY

By S. F. SNIESZKO, *Bacteriologist*, and G. L. BULLOCK, *Fishery Aid*

During the past 2 years, the Microbiological Laboratory, at Leetown, W. Va.,[1] has received an increasing number of reports of outbreaks of furunculosis that failed to respond to treatment with sulfonamides. The problems resulting from furunculosis, a widespread, bacterial disease of salmonids, and control of the disease have been reviewed in detail by McCraw (1952) and Snieszko (1954a). The latter investigator recommended the use of antibiotics for diseased fish that failed to respond to treatment with sulfonamides. In some of the recent incidences of furunculosis at the Leetown station the disease did not respond to sulfonamide therapy, but control was effected with chloramphenicol. Failure of sulfonamides to control furunculosis suggested that the causal organism, *Aeromonas salmonicida*, was resistant to the drugs used.

Several methods are widely used in the routine determination in vitro of the sensitivity of bacteria to sulfonamides and antibiotics. The results obtained with antibiotics are generally accepted as reliable aids in the selection of the most promising treatment. Such, however, is not the case with sulfonamides, and "The relation of *in vitro* sensitivity test to clinical effectiveness is still controversial" (Burdette, Plank, and Clapper, 1955).

The experiments presented in this paper followed three lines of endeavor: (1) A comparison of the effectiveness of sulfonamide and antibiotic therapy in strains of trout which were either susceptible or resistant to furunculosis. (2) Comparison of the therapy of furunculosis with sulfonamides and chloramphenicol in trout which were suffering from furunculosis caused by sulfonamide-resistant, but chloramphenicol-sensitive, strains of *A. salmonicida*. (3) The development of a standard method for laboratory and field use of determining in vitro the sensitivity of *A. salmonicida* to sulfonamides and antibiotics.

Many factors are capable of affecting in-vitro tests of microbial sensitivity to sulfonamides; therefore, correct interpretation and application of the results of such tests will be reliable if a proved method of obtaining reproducible results is used.

As the result of the findings presented in this study and practical experience gained in the treatment of this disease in trout hatcheries over the entire country, a revised procedure has been evolved and described for the treatment of this disease.

The authors wish to express their thanks to Dr. K. E. Wolf for his assistance during the preparation of the manuscript and for furnishing the strains of *A. salmonicida* isolated in Iowa, Utah, Minnesota, and Wisconsin; and to Dr. R. E. Lennon for supplying the fingerling brook trout from Erwin, Tenn.

MATERIALS AND METHODS

Treatment with sulfonamides and chloramphenicol

Experiments on the therapy of furunculosis were carried out with two strains of fingerling brook trout (*Salvelinus fontinalis*) and a strain of brown trout (*Salmo trutta*). A strain of brook trout from Bellefonte, Pa., was selected because it was known to be resistant to furunculosis (Wolf 1954; Snieszko 1954b), and a strain from Erwin, Tenn., was used because it was suspected to be susceptible. The brown trout were from Cortland, N. Y. In general, brown trout are known to be more resistant to furunculosis than are most strains of brook trout (McCraw 1952; Wales and Berrian, 1937).

Equal weights of the trout were distributed among stainless steel troughs in which the water was maintained at temperatures of 12° to 13° C. (54°–55° F.). Infection of the fish was initiated by adding fresh cultures of *Aeromonas salmonicida* to the diet, and treatment was started when the first mortality due to furunculosis occurred.

[1] Post Office, Kearneysville, W. Va.
Note—Approved for publication, March 12, 1957. Fishery Bulletin 125.

Sulfonamides were administered at the rate of 200 milligrams per kilogram of trout per day, or 9 grams per 100 pounds of fish. Chloramphenicol[2] was given at the rate of 75 milligrams of pure antibiotic activity per kilogram of fish per day, or 3.4 grams per 100 pounds of fish. Treatment was continued to the end of the observations recorded in tables 1 to 3. Drugs were mixed with the quantities of food calculated from the Cortland Hatchery feeding charts (Deuel et al., 1952). A daily record of mortality was kept and most of the dead trout were examined bacteriologically.

Drug sensitivity tests in vitro

The drug sensitivity tests were performed with 84 cultures isolated from the experiment on therapy here described or procured during the past years from other experiments or received from other trout hatcheries (table 4). The routine qualitative sulfonamide and antibiotic sensitivity tests were run with commercial multiple sensitivity paper disks[3] on solid bacteriological media. Sensitivity to sulfonamides has also been determined in a quantitative manner by preparation of serial dilutions of these drugs in liquid bacteriological media.

The following solid bacteriological media were used with the paper disks:

1. Leetown standard furunculosis medium having the following composition:

Tryptic digest of casein	10 grams.
Yeast extract	5 grams.
Sodium chloride	2.5 grams.
Agar	15.0 grams.
Water	1 liter.
pH	6.8–7.0.

1A. The same, but buffered with 0.5 percent of secondary sodium phosphate and adjusted to pH 8.0.

2. Sensitivity Test medium C (Case Laboratories).

2A. The same, but buffered and adjusted as No. 2.

3. Mueller Hinton medium (Bacto).

4. Trypticase Soy agar (Baltimore Biological Laboratory).

For the inoculation of the solid media, 24-hour old cultures in nutrient broth were used. These were streaked with cotton swabs on 10-cm. diameter petri plates, each containing 20 ml. of medium, and the disks were placed in position.

The quantitative sensitivity tests were run in liquid media. In order to select the most satisfactory substrate, the following media were tested:

Leetown standard furunculosis medium without agar.

Vitamin-free casamino acids (Difco), 1 percent.

Tryptic digest of casein, 1 percent.

Nutrient broth (Difco).

Nutrient broth (Difco) buffered with 0.5 percent of secondary sodium phosphate and adjusted to desired pH.

Proteose peptone No. 3 (Difco), 1 percent.

N-Z-Case (Sheffield Chemical Co.), 1 percent.

Nutrient broth buffered with sodium phosphate was selected as most promising for our purposes.

Because most of the sulfonamides are only slightly soluble in water, stock solutions of sodium salts were prepared by dissolving a known quantity of sulfonamide in diluted sodium hydroxide. Excess alkali over that needed for keeping sulfonamides in solution was neutralized with hydrochloric acid. Stock solutions were sterilized by filtration, stored under refrigeration, and used within a week. Serial dilutions of 1:25 or 1:5 stock solutions were carried out in 5 ml. of media. The first tube contained double-strength medium and after the serial dilutions were made 5 ml. were discarded from the last tube. Inocula consisted of 0.1 ml. of a 24-hour-old culture diluted 1:5 with sterile water. All cultures were incubated at 20° C. and results were read as soon as abundant growth developed, usually within 24 to 48 hours.

RESULTS

Treatment with sulfonamides and chloramphenicol

The response to treatment with sulfonamides and chloramphenicol of experimental furunculosis in two strains of brook trout and one strain of brown trout is presented in tables 1 to 3. Sulfonamide treatment of experimental furunculosis in the Erwin brook trout failed completely (table 1). Losses were high because the trout were susceptible to furunculosis and the strain of *A. salmonicida* used to produce the disease developed resistance to sulfonamides. When the experiment

[2] Racemic chloromycetin, Parke, Davis & Co., was used at a double rate, because the manufacturer advised that it has about 50 percent the therapeutic value of the d-rotatory isomer.

[3] Multidisks, Case Laboratories, Chicago, Ill.

was repeated with the same strains of trout and bacteria, but with treatment with chloramphenicol, the response was rapid and favorable.

A similar experiment was run concurrently with a strain of brook trout from Bellefonte, Pa., that is known to be resistant to furunculosis. The fish failed to respond to sulfonamide treatment and bacteria isolated from dead trout were found also to be mostly resistant to sulfonamides. Trout surviving the sulfonamide treatment were given chloramphenicol and losses stopped within the first 5-day period (table 2). The overall fish losses were much lower in the resistant Bellefonte strain than in the susceptible Erwin strain.

The results with brown trout (table 3) were very similar to those obtained with Bellefonte brook trout since they are known to be more resistant to furunculosis than most of the strains of brook trout.

TABLE 1.—*Mortality of a strain of eastern brook trout (Erwin, Tenn.), susceptible to furunculosis, following treatment with sulfonamides and chloramphenicol*

A. TREATMENT WITH SULFONAMIDES [1]

[Initial weight of fish per lot (trough), 1,500 gm.; mean weight per fish, 19.8 gm.; mean number of fish per lot (trough), 76; fish infected orally July 29 and Aug. 1, 1955, and treatment started Aug. 8; mortality expressed as percent of fish at beginning of each period]

Period	Mortality under treatment with—				Mortality in controls	
	Sulfamerazine		Sulfisoxazole (Gantrisin [2])			
	Lot A	Lot B	Lot A	Lot B	Lot A	Lot B
Aug. 7–11	7.8	6.5	6.8	2.6	9.1	12.0
Aug. 12–16	12.6	16.6	22.0	39.1	26.0	55.0
Aug. 17–21	37.0	63.3	49.0	75.5	48.0	67.0
Aug. 22–24	33.3	82.0	48.1	63.6	41.0	80.0
Total mortality	66.2	95.0	81.0	94.7	79.2	97.3
Average, total mortality	80.6		87.9		88.2	

B. TREATMENT WITH CHLORAMPHENICOL

[Initial weight of fish per lot (trough), 1,500 gm.; mean weight per fish, 30 gm.; mean number of fish per lot (trough), 50; fish infected orally Sept. 8 and 9, 1955, and treatment started Sept. 16; mortality expressed as percent of fish at beginning of each period]

Period	Mortality under treatment in—		Mortality in controls	
	Lot A	Lot B	Lot A	Lot B
Sept. 16–20	4.1	14.0	9.1	19.1
Sept. 21–25	11.0	16.0	32.0	45.0
Sept. 26–30	0	0	38.2	67.0
Oct. 1–5	0	0	50.0	100.0
Total mortality	14.2	27.5	87.2	100.0
Average, total mortality	20.8		93.5	

[1] Strains of *A. salmonicida* isolated from dead trout were resistant to sulfonamides.

[2] Gantrisin used in these studies was supplied free of cost by Hoffmann La Roche, Nutley, N. J.

TABLE 2.—*Mortality of a strain of eastern brook trout (Bellefonte, Pa.), resistant to furunculosis, following treatment with sulfonamides and chloramphenicol*

A. TREATMENT WITH SULFONAMIDES [1]

[Initial weight of fish per lot (trough), 1,500 gm.; mean weight per fish, 27.6 gm.; mean number of fish per lot (trough), 54; fish infected orally July 29 and Aug. 1, 1955, and treatment started Aug. 19; mortality expressed as percent of fish at beginning of each period]

Period	Mortality under treatment with—				Mortality in controls	
	Sulfamerazine		Sulfisoxazole			
	Lot A	Lot B	Lot C	Lot D	Lot E	Lot F
Aug. 17–21	0	1.8	0	4.0	3.6	0
Aug. 22–26	0	7.5	1.6	0	7.5	4.0
Aug. 27–31	3.5	4.0	1.6	0	8.1	4.1
Sept. 1–5	1.8	10.6	10.1	0	4.4	0
Total mortality	5.3	22.2	13.1	4.0	21.8	8.0
Average, total mortality	13.7		8.5		15.0	

B. TREATMENT WITH CHLORAMPHENICOL

[Fish surviving treatment with sulfonamides (2-A) used in this experiment; weight of fish per lot (trough), 2,000 gm.; mean weight per fish, 43 gm.; mean number of fish per lot (trough), 46; no additional infection, with treatment started Sept. 7, 1955; mortality expressed as percent of fish at beginning of each period]

Period	Mortality under treatment in—			Mortality in controls		
	Lot A	Lot C	Lot E	Lot B	Lot D	Lot F
Sept. 6–10	9.4	7.5	0	12.0	0	4.3
Sept. 11–15	0	0	0	10.8	0	0
Sept. 16–20	0	0	0	3.0	2.1	0
Sept. 21–25	0	0	0	3.1	0	2.2
Sept. 26–30	0	0	0	0	2.1	2.3
Total mortality	9.4	7.5	0	26.1	4.2	8.7
Average, total mortality	5.7			10.8		

[1] Strains of *A. salmonicida* isolated from dead fish were resistant to sulfonamides.

TABLE 3.—*Mortality of brown trout following treatment with sulfonamides and chloramphenicol*

A. TREATMENT WITH SULFONAMIDES [1]

[Initial weight of fish per lot (trough), 1,500 gm.; mean weight per fish, 22.0 gm.; mean number of fish per lot (trough), 68; fish infected orally July 29 and Aug. 1, 1955, and treatment started Aug. 12; mortality expressed as percent of fish at beginning of each period]

Period	Mortality under treatment with				Mortality in controls	
	Sulfamerazine with sulfaguanidine		Sulfisoxazole			
	Lot A	Lot B	Lot C	Lot D	Lot E	Lot F
Aug. 12–16	3.1	3.0	1.3	1.7	1.4	6.6
Aug. 17–21	0	1.5	0	3.4	0	5.2
Aug. 22–26	4.8	3.1	2.7	0	1.4	13.0
Aug. 27–31	10.0	8.0	5.5	7.1	8.7	4.2
Sept. 1–5	5.6	10.3	8.8	3.8	4.8	2.2
Total mortality	22.0	25.7	17.3	17.5	17.8	29.0

[1] Strains of *A. salmonicida* isolated from dead fish were resistant to sulfonamides.

TABLE 3.—*Mortality of brown trout following treatment with sulfonamides and chloramphenicol*—Con.

B. TREATMENT WITH CHLORAMPHENICOL

[Fish surviving treatment with sulfonamides (3-A) used in this experiment; weight of fish per lot (trough), 1,500 gm.; mean weight per fish, 31 gm.; mean number of fish per lot (trough), 48; no additional infection, with treatment started Sept. 7, 1955; mortality expressed as percent of fish at beginning of each period]

Period	Mortality under treatment in—			Mortality in controls		
	Lot A	Lot C	Lot E	Lot B	Lot D	Lot F
Sept. 6–10	10.0	1.6	3.3	13.4	0	6.8
Sept. 11–15	2.2	0	0	8.9	6.1	7.3
Sept. 16–20	0	0	0	0	6.4	7.9
Sept. 21–25	0	1.6	0	0	0	5.7
Sept. 26–30	0	1.6	0	0	2.3	6.1
Total mortality	12.0	4.8	3.3	21.0	14.0	29.5
Average, total mortality	6.7			21.3		

While the results obtained in the treatment of furunculosis caused by sulfonamide-resistant but chloramphenicol-sensitive strains of *A. salmonicida* are of considerable interest in their own merit, they are presented in this paper chiefly as background information for the history of the majority of the strains of this bacterium which were used in the in-vitro sensitivity tests. Summary information on the cultures used for testing o the drug sensitivity is presented in table 4.

TABLE 4.—*Sensitivity of 84 strains of* A. salmonicida *to sulfonamides tested with sodium sulfadiazine in buffered nutrient broth*

Date and source of culture	Place of isolation	Number of strains that were—			
		Sensitive [1]	Intermediate	Resistant [2]	Total
1955:					
Brook trout:					
Not treated	Leetown	0	2	12	14
Treated with sulfonamides.	do	0	1	9	10
Treated with chloramphenicol.	do	0	0	5	5
Brown trout:					
Not treated	do	1	5	1	7
Treated with several sulfonamides.	do	2	0	9	11
Total		3	8	36	47
1956:					
Brown trout: Not treated	Leetown	13	2	2	17
Rainbow trout: Not treated	Iowa	2	0	0	2
Do	Utah	0	0	2	2
Brown and brook trout: Treatment (?).	Minnesota	4	0	0	4
Brook trout: Treatment (?)	Wisconsin	3	0	0	3
Total		22	2	4	28
Before 1955:					
Miscellaneous stock cultures stored in lyophilized form since 1952.	Leetown	4	2	3	9
Grand total		29	12	43	84

[1] Cultures in which there was no growth within 24 hours in buffered nutrient broth having a pH of 8.0 and containing sodium sulfadiazine in the concentration of 1:1,600.

[2] Cultures in which there was undiminished growth in nutrient broth containing sodium sulfadiazine in the concentration of 1:100.

Sensitivity tests with antibiotics

Tests of the sensitivity of *Aeromonas salmonicid* to various antibiotics were performed with 4 strains of the pathogen that were isolated durin the summer of 1955 from fingerling brook trou known to be either susceptible or resistant t furunculosis and from fingerling brown trout Trout from which the bacteria were isolated ha been treated with sulfonamides or chloramphe nicol, or had served as controls (table 4). Cul tures of these strains were grown on the Leetow standard furunculosis medium on which 6-tippe sensitivity disks containing various antibiotic had been placed (table 5 and fig. 1). All strain of *A. salmonicida* that were tested were found t be uniform in their response to disks containin antibiotics. No correlation between the sensitiv ity to antibiotics and sulfonamides was apparent

TABLE 5.—*Sensitivity of* Aeromonas salmonicida *to selecte antibiotics as determined with Multidisks*

Antibiotic	Quantity of antibiotic in disk	Width of zones of growth inhibition (mm.)
Chloromycetin [1]	10 μg	14–1
Terramycin [2]	do	12–1
Tetracycline	do	10–1
Aureomycin [3]	do	10–1
Erythromycin	do	5–
Streptomycin	do	4–
Neomycin	do	1–
Penicillin	1.5 units	0–
Carbomycin	5 μg	
Bacitracin	5 units	
Polymyxin B	10 units	
Viomycin	10 μg	
Furadantin [4]	50 μg	8–1

[1] Chloramphenicol.
[2] Oxytetracycline.
[3] Chlortetracycline.
[4] Nitrofurantoin; related to furacin, which has possibilities in the treat ment of furunculosis (Gutsell 1948).

Sensitivity tests with sulfonamides

Tests were run in liquid and solid media. Th first liquid substrate tested was the standar furunculosis medium containing serial dilutions o sodium sulfamerazine. The pH of the medium wa 6.8 to 7.0 and the sulfonamide concentration ranged from 1:100 to 1:10,000. Thirty resistan and 10 sensitive strains of *A. salmonicida* wer used. The results were not satisfactory becaus sensitive strains of *A. salmonicida* grew as well a the resistant strains. Crystals of sulfamerazin appeared in test tubes with the higher drug con centrations. Therefore, other liquid media suc



FIGURE 1.—Sensitivity of *Aeromonas salmonicida* in vitro to various antibiotics as determined with three media: Trypticase Soy agar (TSA), Mueller Hinton agar (MH), and standard furunculosis agar (#4). The antibiotic and quantity used on Multidisks, reading clockwise, were as follows: Upper: Tetracycline (TE), 10μg.; erythromycin (E), 10 μg.; terramycin (T), 10 μg.; carbomycin (CA), 5 μg.; bacitracin (B), 5 units; aureomycin (A), 10 μg. Lower: Viomycin (V), 10 μg.; chloromycetin (C), 10 μg.; polymyxin B_1 (PB), 10 units; penicillin (P), 1.5 units; streptomycin (dihydro) (S), 10 μg.; neomycin (N), 10 μg. The presence of a clear zone indicates sensitivity.

as proteose peptone #3, vitamin-free casamino acids, nutrient broth, and N-Z-Case were tested. Surprisingly, the most promising results were obtained with nutrient broth, and this medium was used in all subsequent experiments unless otherwise indicated. Since the sodium salts of sulfonamides give alkaline solutions and their solubility and ionization decreases with decreasing alkalinity of the medium, nutrient broth was buffered by adding 0.5 percent of secondary sodium phosphate and the pH of the medium adjusted as required. In the unbuffered nutrient broth (original pH, 6.8) sodium sulfamerazine precipitated if added in concentration up to 1:250. In such a medium the sulfa-sensitive and sulfa-resistant strains of *A. salmonicida* grew equally well.

Since Griffin et al. (1953) found that *A. salmonicida* has a wide pH tolerance, the buffered nutrient broth was adjusted to pH 7.0, 7.5, 8.0, 8.5, and 9.0. The more alkaline the medium, the more striking were the differences noted between the sulfa-sensitive and sulfa-resistant strains. The all-round best results were obtained at pH 8.0 to 8.5; therefore, all subsequent tests were run within this pH range unless otherwise specified.

According to Northey (1948), "Frequently it is found that once an organism has become resistant to one of the potent sulfonamide drugs it is resistant to all the commonly used sulfa drugs." Therefore it should be advantageous to select for a routine in-vitro test with *A. salmonicida* a sulfonamide, or sulfonamides, which would be the most convenient to work with. With this in mind, several commonly used sulfonamides were tested with the two resistant and two sensitive strains of *A. salmonicida*. Sodium salts prepared in the laboratory were serially diluted in the buffered nutrient broth adjusted to pH 8.0 to 8.5. The results are presented in table 6. Results for sulfaguanidine are omitted from the table because a soluble sodium salt could not be prepared within the desired pH range.

On the basis of these results, all available strains (84) of *A. salmonicida* were tested with sulfadiazine, though sulfamerazine or sulfisoxazole could be used equally well. Results are presented in table 4. It is evident that the largest number of resistant cultures were isolated at Leetown in 1955 from treated or control trout during a period when treatments with sulfonamides were being carried out. Cultures isolated at Leetown in 1956 before any treatments started were predominantly sensitive. There were some intermediate strains which produced a scanty growth in the medium with sodium sulfadiazine.

The highest dilutions of sulfonamides arresting growth of *A. salmonicida* in a buffered nutrient

TABLE 6.—*Growth of sulfonamide-resistant and sulfonamide-sensitive strains of* Aeromonas salmonicida *in buffered, nutri broth with various sulfonamides*

[Question mark (?) indicates reliable reading could not be made because of precipitate in medium]

Strain number and sulfonamide dilution	Sulfonamides (sodium salt)							
	Sulfadiazine	Sulfamerazine	Sulfamethazine	Sulfathiazole	Sulfisoxazole	Sulfaquinoxaline	Sulfathialidine	Sulfanilamide
Sensitive #1:								
1:50	?	?	?	?	−	?	?	?
1:100	−	−	?	?	−	?	+++	?
1:200	−	−	+	−	−	?	+++	?
1:400	−	−	+++	+	−	−	+++	?
1:800	−	−	+++	+++	−	−	+++	+
1:1600	−	−	+++	+++	−	−	+++	++
Control	+++	+++	+++	+++	+++	+++	+++	+++
Sensitive #2:								
1:50	?	?	?	?	−	−	?	?
1:100	−	−	?	?	−	+	+++	?
1:200	−	−	−	−	−	+	+++	?
1:400	−	−	−	+	−	+	+++	?
1:800	−	−	−	+	−	+	+++	+
1:1600	−	−	−	+	−	+	+++	++
Control	+++	+++	+++	+++	+++	+++	+++	+++
Resistant #1:								
1:50	?	?	?	?	−	−	?	?
1:100	++	−	?	?	−	−	+++	?
1:200	+++	+++	+	−	+	−	+++	?
1:400	+++	+++	+++	+	+++	+	+++	?
1:800	+++	+++	+++	+++	+++	+++	+++	+
1:1600	+++	+++	+++	+++	+++	+++	+++	++
Control	+++	+++	+++	+++	+++	+++	+++	+++
Resistant #2:								
1:50	?	?	?	?	−	−	?	?
1:100	+++	−	+	?	+	−	+++	?
1:200	+++	+++	+++	−	+++	+	+++	?
1:400	+++	+++	+++	+	+++	+++	+++	?
1:800	+++	+++	+++	+++	+++	+++	+++	+
1:1600	+++	+++	+++	+++	+++	+++	+++	++
Control	+++	+++	+++	+++	+++	+++	+++	+++

broth were determined with laboratory-prepared sodium salts of sulfadiazine, sulfamerazine, sulfamethazine, and sulfisoxazole. Ten strains of sulfonamide-sensitive and 10 resistant cultures were used. Serial dilutions of sulfonamides ranged from 1:100 to 1:102,400. Control test tubes contained medium without sulfonamides. After 24 hours of incubation the resistant strains grew well in a sulfonamide concentration of 1:100 (1:200 with sulfamerazine). Growth of the sensitive strains was absent in the following concentrations and in all lower dilutions:

Sulfadiazine	1:12,800
Sulfamerazine	1:6,400
Sulfamethazine	1:6,400
Sulfisoxazole	1:6,400

The amount of growth was much reduced up to the highest dilution of the sulfonamides tested (1:102,400, or 10 μg. per ml.). After 48 hours the results differed very little except in the medium with sulfamethazine which showed abundant growth up to 1:200 and scanty at 1:100 (the highest concentration tested).

The final series of experiments was conducted for the purpose of selecting a reliable method of determining the drug sensitivity of *A. salmonicida* under either laboratory or field conditions.

In a preliminary test, the optimum pH rang for the appearance of growth inhibition zones o an agar medium was determined. Filter pap disks (6 mm. in diameter) [4] were impregnate with 1 mg. of filter-sterilized sodium sulfadiazi dissolved in 0.02 ml. of water and air dried 37° C. Standard furunculosis medium (buffere adjusted to pH 7.0, 7.5, 8.0, 8.5, and 9.0, a sterilized) was distributed in 20-ml. quantities 10 petri plates (10-cm. in diameter), the mediu was inoculated by swabbing the surface with 24-hour culture of *A. salmonicida*, and dis were positioned. Results were recorded after 2 and 48 hours of incubation at 20° C. Zon were distinctly visible after 24 hours, but t width of the zones could be more convenient determined after 48 hours. The widths of t zones with sulfonamide-sensitive strains at vario pH levels were as follows:

pH 7.0	11 mm.
pH 7.5–8.0	15 mm.
pH 8.5	16 mm.
pH 9.0	17 mm.

There were no clear zones with sulfa-resista strains. The best results, based on the width

[4] No. 740–E., C. Schleicher & Schuell Co.

he zones and abundance and rapidity of growth, vas in the medium with pH 8.0.

As a result of this preliminary test to determine he optimum pH range, another experiment was carried out with 6-tipped commercial multiple disks and ingle disks prepared in the laboratory. On standrd furunculosis medium, zones of partial growth nhibition appeared around disks with 1 milligram f sodium sulfisoxazole, 1.0 and 0.5 milligram of odium sulfadiazine, and 0.1 milligram of thiolfil.[5] No zones appeared around disks containing .1 milligram of sulfisoxazole, sulfadiazine, elkosin, ulfamerazine, or triple sulfa.[6] The same test vas repeated with (1) Case Laboratory Sensitivity Test medium "C," as prepared by the manufacturer, and (2) the same medium buffered and lkalized to pH 8.0. Results are presented in able 7.

Growth of *A. salmonicida* was strikingly less bundant on nonbuffered "Case" medium, probbly because of the presence of dextrose. To test he effect of dextrose, a standard furunculosis nedium buffered and adjusted to pH 8.0 was enriched with 0.5 percent of this sugar. On such a nedium and with multiple disks containing as nuch as 0.5 milligram of sulfonamides per tip, no lear zones appeared with sensitive cultures.

TABLE 7.—*Sensitivity of sulfa-sensitive strains of* A. salmonicida *to sulfonamides cultured on Sensitivity Test medium "C" (Case)*

[L=disks prepared at Leetown; PZ=zones of partial growth inhibition]

Quantity of sulfonamide used	Average zone width (in mm.) on medium—	
	Not modified	Buffered and alkalized
odium sulfadiazine:		
0.1 milligram	1.3	2.8
0.1 milligram (L)	1.8(PZ)	8.4(PZ)
0.5 milligram (L)	6.9(PZ)	15.0
1.0 milligram (L)	9.7(PZ)	17.5
odium sulfisoxazole:		
0.1 milligram	6.8(PZ)	6.1
1.0 milligram (L)	13.5(PZ)	20.3
ulfamerazine: 0.1 milligram	0	3.6
hiosulfil: 0.1 milligram	4.5	3.6
riple sulfa: 0.1 milligram	0	2.1

A similar test was performed with commercial Mueller Hinton and Trypticase Soy agar media and 7-tipped Multidisks containing 0.1 or 0.5 milligram of sulfonamides per disk. The width of he zones could not be recorded because the clear zones were so wide that practically no growth appeared on the 10-cm. petri plates. Therefore the test was repeated with single disks, prepared at the Leetown laboratory, which contained 0.1, 0.25, and 0.5 milligram of sodium sulfadiazine.

The results presented in table 8 show that the lack of growth on 10-cm. petri plates using multiple disks was due to the great width of zones produced by disks containing more than 0.1 milligram of sulfonamide. Therefore use of multiple disks with higher concentrations of sulfonamides and with Mueller Hinton and Trypticase Soy agar media in 10-cm. petri plates is not advisable.

TABLE 8.—*Determining sulfonamide sensitivity of* A. salmonicida *using two media and single disks*

Quantity of sodium sulfadiazine per disk	Width of clear zone (mm.) on—	
	Mueller Hinton medium (Difco)	Trypticase Soy agar (BBL)
	Mm.	*Mm.*
0.1 milligram	12	7–8
0.25 milligram	13–15	10
0.5 milligram	15–16	12–13

The sulfonamide-sensitivity test in which filter paper disks containing drugs were employed could be much more conveniently performed under field conditions if petri plates were replaced by test tubes. Therefore, the buffered and alkalized standard furunculosis medium, Mueller Hinton medium (Difco), and Trypticase Soy agar (BBL) were used as agar slants. Media were inoculated by making longitudinal streaks with a loop containing a suspension of *A. salmonicida*, and disks were deposited in the center of the slant. Excellent clear zones appeared (fig. 2) with sulfa-sensitive strains.

Another approach to the performance of the sulfonamide-sensitivity test under simulated field conditions was made with agar media containing sodium sulfadiazine added to the buffered and alkalized standard furunculosis medium. These solutions were sterilized by filtration or were incorporated in the medium before sterilization in the autoclave. In all cases, clear differences were noticed in the growth of sulfonamide-sensitive and sulfonamide-resistant strains of *A. salmonicida*. The optimum concentrations of sodium sulfadiazine in the medium were 1:500 and 1:1,000.

It is expected that extensive field trials based on the results presented here will permit selection

[5] Ayerst Laboratories.
[6] A mixture of sulfamerazine, sulfamethazine, and sulfadiazine.



FIGURE 2.—Sulfonamide-sensitivity test performed with a sensitive culture of *A. salmonicida* using Trypticase Soy agar (left) and Mueller Hinton agar (right). Laboratory-prepared disks contained 0.1, 0.25, and 0.5 milligram of sodium sulfadiazine.

of the most convenient method for the determination of the sensitivity of fish-pathogenic bacteria to sulfonamides, antibiotics, and perhaps, to other therapeutic agents.

DISCUSSION

Micro-organisms appear to be capable of developing resistance to the majority of drugs which are employed clinically. In view of the cross-resistance shown between different groups of drugs, it is obviously important that the clinician should have information as to which groups of drugs may be usefully tried in infections with resistant strains (Work and Work, 1948).

It has been found, as could be expected, that with the wide use of sulfonamides for treatment of fish diseases, there is an increasing number of outbreaks of furunculosis which are refractory to treatment with these drugs. Since "once resistance is highly developed it is apparently permanent" (Northey 1948), it is increasingly important to have means of rapid and reliable determination of such resistance.

In this paper, observations are described on the treatment of experimentally induced furunculosis employing strains of trout which are susceptible or resistant to this disease. The outbreaks were caused by strains of the pathogen which were found to be either sensitive or resistant to sulfonamides. When it was found that the disease was refracto to treatment with sulfonamides, chloramphenic was used with good results (tables 1 to 3). Ox tetracycline (terramycin), which is also effecti in the treatment of furunculosis and ulcer disea (Snieszko et al., 1952), was not used in this inves gation because strains of *A. salmonicida* resista to this antibiotic have recently been isolated.

It is evident from the results described he that in addition to chloramphenicol and oxytet cycline other antibiotics and sulfonamides arr the growth of *A. salmonicida* in vitro. The resu of these tests cannot be applied indiscriminate for therapeutic use. While some of the sulfo amides gave promising results in vitro, their u in the treatment of fish diseases may be limited their toxicity to some species of fish or by t arresting of the fish's growth during treatme Other drugs may not be absorbed from the i testinal tract and are therefore useless in system infections by micro-organisms sensitive to the in vitro (Johnson and Brice, 1953; Snieszko a Wood, 1955; Snieszko and Friddle, 1951; Sniesz and Griffin, 1955). It is interesting that chlo tetracycline (aureomycin) which has been fou to be effective in vitro against *A. salmonicida* a *Hemophilus piscium* is entirely ineffective in t treatment of trout suffering from furunculosis a ulcer disease (Snieszko, Griffin, and Friddle, 195 Therefore, the in-vitro tests, while very convenie because of the speed and ease with which they c be performed, must be supplemented by exa treatment trials with several species of fish before the therapeutic value of drugs can determined.

The greatest value of the in-vitro test is in t determination of the acquired drug resistance the pathogen concerned. This can be explained be by using an example. It is known that furunc losis usually can be effectively treated with su fonamides. It has also been shown in this pap that the growth of sensitive strains of *A. salmon cida* can be arrested in vitro with these drug Therefore, if it is found that a particular strain *A. salmonicida* is resistant to sulfonamides in vitr an outbreak of furunculosis caused by that stra will most likely be refractory to a treatment wi sulfonamides. It does not mean, however, th if the organism is found to be sensitive to a certa drug in vitro, the treatment with that drug of fi infected with such an organism will also be effe

ive. The experience with chlortetracycline previously cited may serve as an example. It is well o remember the limitations of the in-vitro test or a correct interpretation of the results.

The results of our studies show that in the in-vitro est sulfadiazine, sulfamerazine, and sulfisoxazole, nd probably other sulfonamides, can be used qually well to determine sulfonamide sensitivity f *A. salmonicida*. In the therapeutic and feeding xperiments, however, sulfadiazine has been found o be more toxic to the fish than has sulfamerazine, nd sulfisoxazole has been found to be free from ny detectable untoward effects even to brown rout (Snieszko and Wood, 1955). Unfortunately, he experiment on the use of sulfisoxazole for the reatment of furunculosis was unsuccessful because he strains of *A. salmonicida* isolated from the utbreak were resistant to sulfonamides [7] (tables to 3).

Testing of the sensitivity of *A. salmonicida* to ulfonamides and antibiotics can be carried out n several ways. The most practical method eems to be the use of drug-containing, filter-paper isks. Selection of a proper medium is very important when testing the sensitivity of the organism to sulfonamides. Trypticase Soy agar, special urunculosis agar medium buffered and alkalized, nd Mueller Hinton agar, are especially recommended. In the first two media, the brown pigment so characteristic of *A. salmonicida* is produced within 48 hours and the paraphenylenediamine test of Griffin (1952) can be performed as oon as growth is visible to the unaided eye. With he first two media, disks should contain 0.5 milligram of the drug; with Mueller Hinton medium, nly 0.1 milligram. The test can be equally well erformed using either petri plates or agar slants—he latter are more suitable for fieldwork. Also nedia in which sulfonamides have been incorporated may be used provided that the control medium is free of sulfonamides and that both sulfonamide-resistant and sulfonamide-sensitive strains of *A. salmonicida* are available for controls.

RECOMMENDATIONS

On the basis of the results reported in this paper and past experience in the treatment of furunculosis, we wish to make the following recommendations:

1. As soon as a disease suspected to be furunculosis breaks out, start treatment with sulfonamides. Sulfamerazine is still considered the drug of choice, but sulfamethazine or a combination of sulfamerazine with sulfaguanidine can also be used (Snieszko 1954a). Sulfisoxazole is a drug of great promise, but it still has to be evaluated experimentally.

2. The diagnosis of furunculosis should be confirmed as soon as possible by bacteriological examination. This should include determination in vitro of the sensitivity of the pathogen to the drug.

3. The dosage of sulfonamides is 8 to 10 grams per 100 pounds of fish per day. Treatment should last for 10 to 20 days and should result in a complete stoppage of mortalities due to furunculosis. Recurrences are particularly likely in disease-susceptible strains of trout.

4. Sulfonamides should never be used at lower levels or treatments repeated at short intervals. Such practice is the surest way to produce sulfa-resistant strains of the pathogen.

5. If the response to the treatment with sulfonamides is not rapid, if recent experience has shown that treatment with sulfonamides is not effective, or if the results of laboratory examination show that the organism is sulfa-resistant but sensitive to oxytetracycline or chloramphenicol, either antibiotic should be used at a rate of 2.5 to 3.0 grams (of the antibiotic activity) per 100 pounds of fish per day. Chloramphenicol is somewhat better than oxytetracycline.

6. One should never rely on any drug for a permanent control of furunculosis, or of any other disease. There is no drug in existence which would permit elimination of any animal or plant disease. The best a drug can do is to reduce losses temporarily. Long-lasting control of furunculosis, or any other infectious disease, is only possible by elimination of the source of infection, good sanitation, and introduction or development by selective breeding, of a strain of fish with greater disease resistance. Good hatchery practices, avoidance of crowding, and balanced nutrition are very important factors in keeping fish healthy.

SUMMARY

Therapeutic studies were performed with three sulfonamides and one antibiotic (chloramphenicol) with two strains of fingerling brook trout—one

[7] One experiment of this series with sulfisoxazole was performed at Leetown in 1955 by Bo Svenonius, a visiting fishery biologist from Sweden.

susceptible to furunculosis; the other resistant to this disease—and one strain of brown trout. Since it was found that the disease was caused by a sulfa-resistant type of the pathogen, only the treatment with the antibiotic was effective.

Testing for drug resistance has shown that many strains of *A. salmonicida* were resistant to sulfonamides, but sensitive to antibiotics. Therefore, studies were made in order to develop a rapid and reliable field method for the determination of drug sensitivity of *A. salmonicida*.

Of all media tested, only the Trypticase Soy agar and the modified Leetown standard furunculosis medium can be used for the determination of sensitivity of *A. salmonicida* to sulfonamides and antibiotics and at the same time for the rapid presumptive test of Griffin. Mueller Hinton agar is excellent for the determination of drug sensitivity, but not for the presumptive test of Griffin involving the production of brown pigment.

A method of the determination of drug sensitivity employing media in the form of agar slants and single disks containing the drug is recommended for field use.

A revised and up-to-date method of treatment and control of furunculosis is described.

LITERATURE CITED

Burdette, Robin I., LeRoy E. Plank, and William E. Clapper.
1955. A comparison of four methods of testing sulfonamide sensitivity and the relation to clinical response in urinary tract infections. Antibiotics and Chemotherapy, vol. 5, pp. 392–397.

Deuel, Charles R., David C. Haskell, Donald R. Brockway, and O. R. Kingsbury.
1952. The New York State fish hatchery feeding chart. Fisheries Research Bulletin No. 3, third edition, 61 pp. New York State Conservation Department, Albany, N. Y.

Griffin, Philip J.
1952. A rapid presumptive test for furunculosis in fish. Progressive Fish-Culturist, vol. 14, pp. 74–75. U. S. Fish and Wildlife Service, Washington, D. C.

Griffin, Philip J., S. F. Snieszko, and S. B. Friddle.
1953. A more comprehensive description of *Bacterium salmonicida*. Trans. American Fisheries Soc., vol. 82 (1952), pp. 129–138.

Gutsell, James S.
1948. The value of certain drugs, especially sul drugs, in the treatment of furunculosis in broo trout, *Salvelinus fontinalis*. Trans. American Fish eries Soc., vol. 75 (1945), pp. 186–199.

Johnson, H. E., and R. F. Brice.
1953. Mortality of silver salmon from treatmen with sulfamerazine. Progressive Fish-Culturis vol. 15, pp. 31–32. U. S. Fish and Wildlife Servic Washington, D. C.

McCraw, Bruce M.
1952. Furunculosis of fish. Special Scientific Re port: Fisheries No. 84, 87 pp. U. S. Fish and Wild life Service, Washington, D. C.

Northey, Elmore H.
1948. The sulfonamides and allied compound xxvii and 660 pp. Reinhold Publishing Co., Ne York.

Snieszko, S. F.
1954a. Therapy of bacterial fish diseases. Sym posium, Research on Fish Diseases. Trans. Amer can Fisheries Soc., vol. 83 (1953), pp. 313–330.
1954b. Advances in the studies of infectious fis diseases. Trans. Meeting Southeastern Fish an Game Commissioners, 3 pp. New Orleans.

Snieszko, S. F., and S. B. Friddle.
1951. Tissue levels of various sulfonamides in trou Trans. American Fisheries Soc., vol. 80 (1950), pp 240–250.

Snieszko, S. F., and P. J. Griffin.
1955. Kidney disease in brook trout and its treat ment. Progressive Fish-Culturist, vol. 17, pp 3–13. U. S. Fish and Wildlife Service, Washing ton, D. C.

Snieszko, S. F., P. J. Griffin, and S. B. Friddle.
1952. Antibiotic treatment of ulcer disease and fu runculosis in trout. Trans. Seventh North Amer ican Wildlife Conference, pp. 197–213.

Snieszko, S. F., and E. M. Wood.
1955. The effect of some sulfonamides on the growth of brook trout, brown trout and rainbow trout Trans. American Fisheries Soc., vol. 84 (1954), pp 86–92.

Wales, J. H., and William Berrian.
1937. The relative susceptibility of various strain of trout to furunculosis. California Fish and Game vol. 23, pp. 147–148.

Wolf, Louis E.
1954. Development of disease-resistant strains o fish. Symposium, Research on Fish Diseases Trans. American Fisheries Soc., vol. 83 (1953), pp 342–349.

Work, Thomas S., and Elizabeth Work.
1948. The basis of chemotherapy. xvi and 435 pp Interscience Publishers, New York.

U. S. GOVERNMENT PRINTING OFFICE: 1957 O—433457

INDEX TO VOLUME 57

	Page
Aeromonas salmonicida	558
Aetideidae	355
Age determination of trout in Lake Michigan	1–59
annular markings	11
Agonostomus monticola, fresh-water mullet	415
description of species	417
locality—Bahamas and South Atlantic	416
occurrence of larval and juvenile forms	424
spawning area	424
Aeromonas salmonicida resistant to drugs	555, 564
Alutera ventralis	337
Anderson, William W.: EARLY DEVELOPMENT, SPAWNING, GROWTH, AND OCCURRENCE OF THE SILVER MULLET (*Mugil curema*) ALONG THE SOUTH ATLANTIC COAST OF THE UNITED STATES	397–414
Anderson, William W.: LARVAL FORMS OF THE FRESH-WATER MULLET (*Agonostomus monticola*) FROM THE OPEN OCEAN OFF THE BAHAMAS AND SOUTH ATLANTIC COAST OF THE UNITED STATES	415–425
Annotated bibliography of tuna and tuna fisheries	173–249
Annular markings on lake trout	11
characteristics of lake trout scales, Michigan	11
arenaria, Mya	279–292
artedi, Leucichthys	87
Atlantic round herring, *Etrumeus sadina*	335
Atlantic lizard fish (*Synodus foetens*)	336
abundance in southern New England	336
Atlantic sailfish	139–169
Antibiotics, Sensitivity tests on	555
Auxis thazard	336
Bacteria, notes on	469–496
Isolated from *Gymnodinium brevis* cultures	488
List of control fishes in unialgal cultures	478
Bahamas and South Atlantic Coast fresh-water mullet	415
Bare Lake, Alaska, salmon fecundity study	457
Bedroom Cove, soft clam growth in	279
Bibliography of tunas and tuna fisheries, 1930–1953	173
Bigelow, Henry B., Clyde C. Taylor, and Herbert W. Graham: CLIMATIC TRENDS AND THE DISTRIBUTION OF MARINE ANIMALS IN NEW ENGLAND	293–345
Blue marlin, *Makaira ampla*	169, 532–538
Boothbay Harbor, Maine, Mean water temperatures	344
Boston otter trawlers, Georges Bank	266
Bradyidius Giesbrecht	355–358
Bradyidius arnoldi, new species	355
Brevis, red tide	475–492
Bullock, G. L., and S. F. Snieszko: TREATMENT OF SULFONAMIDE-RESISTANT FURUNCULOSIS IN TROUT AND DETERMINATION OF DRUG SENSITIVITY	555–564
Butterfly ray (*Gymnura altavela*), south of Cape Cod	335
Cable, Louella E.: VALIDITY OF AGE DETERMINATION FROM SCALES, AND GROWTH OF MARKED LAKE MICHIGAN LAKE TROUT	1–39
caerulea, Sardinops, Pacific sardine	427–449
Calanoida	347
Calanoid copepods of families Aetideidae, Euchaetidae, and Stephidae from the Gulf of Mexico	355–363
Carcinides maenas, green crab	340
Carnegie (1929) zooplankton collections, Central Pacific	365
Central Equatorial Pacific, Yellowfin tuna spawning	251–264
Capelin, Pacific Coast, *Mallotus catervarius*	444
catervarius, Mallotus, capelin of Pacific Coast	444
Central Pacific, food of bigeye and yellowfin tuna	64, 82
Central Pacific, Observations on the spearfishes	497–554
Central Pacific, Zooplankton abundance	365
Central Pacific, Zooplankton studies, 1951–54	365–395
Centropristes striatus, sea bass, Corea, Maine	337
cephalus, Mugil	415–421
Charlevoix (Mich.), U.S. Fish Hatchery	2
Chloramphenicol, Treatment with	555
Clam, growth of *Mya arenaria*, soft	279, 292
CLIMATIC TRENDS AND THE DISTRIBUTION OF MARINE ANIMALS IN NEW ENGLAND, by Clyde C. Taylor, Henry B. Bigelow, and Herbert W. Graham	293–345
Clupea pallasi, Pacific herring spawning compared with *sardinops*	443
relative abundance of eggs of herring	451
Collybus drachme, pomfret, food item of tuna	66, 75, 79
COMPARATIVE STUDY OF FOOD OF BIGEYE AND YELLOWFIN TUNA IN THE CENTRAL PACIFIC, by Joseph E. King and Isaac I. Ikehara	61–85
Copepoda calanoida	347–354
Copepods, New calanoid	355
Coregonus (Artedi) Linnaeus, Great Lakes	87
in Europe	87
Coryphaena hippurus, common dolphin, northern range	377
Crab, green (*Carcinides maenas*)	340
curema, Mugil, silver mullet	397–414
Cyprinodon variegatus, experimental	486, 495

Page

Depth zones, Distribution of different species in 274
Development of silver mullet, larval to juvenile stage 402
Dolphin, common (*Coryphaena hippurus*) 377
Drug resistance 556
furunculosis in trout 555–564

EARLY DEVELOPMENT, SPAWNING, GROWTH, AND OCCURRENCE OF THE SILVER MULLET (*Mugil curema*) ALONG THE SOUTH ATLANTIC COAST OF THE UNITED STATES, by William W. Anderson 397–414
Ecological notes on sailfish 167
EFFECTS OF ENVIRONMENT AND HEREDITY ON GROWTH OF THE SOFT CLAM (*Mya arenaria*), by Harlan S. Spear and John B. Glude 279–292
EFFECTS OF UNIALGAL AND BACTERIA-FREE CULTURES OF *Gymnodinium brevis* ON FISH, and notes on related studies with bacteria, by Sammy M. Ray and William B. Wilson 469–496
Elbow Cay, Bahamas, fresh-water mullet collections 415
Estimating abundance of groundfish on Georges Bank 265–278
Etvumeus sadina, herring, northern range 335
Euchaeta paraconcinna, new species 358
Euchaetidae 358–361
Eulachon, *Thaleichthys pacificus*, (Pacific Coast) spawning 443
Euthynnus alletteratus, little tuna, northern range 336
pelamis, striped bonito, northern range 336

FECUNDITY OF NORTH AMERICAN SALMONIDAE, by George A. Rounsefell 451–468
FECUNDITY OF THE PACIFIC SARDINE (*sardinops caerulea*), by John S. MacGregor 427–449
Filefish (*Alutera ventralis*) from Tortugas region 337
Fin regeneration, Lake Michigan trout 7
Fishing of otter trawlers in each depth zone 270
Flavobacterium piscicida, bacterial counts 490, 491
Fleminger, Abraham: NEW CALANOID COPEPODS OF THE FAMILIES AETIDEIDAE, EUCHAETIDAE, AND STEPHIDAE FROM THE GULF OF MEXICO 355–363
Fleminger, Abraham: NEW GENUS AND TWO NEW SPECIES OF THARYBIDAE (*Copepoda calanoida*) FROM THE GULF OF MEXICO, WITH REMARKS ON THE STATUS OF THE FAMILY 347–354
Florida Coast M/V *Theodore N. Gill* collections of mullet 415, 416
Food habits, sailfish 164
Food of yellowfin and bigeye tuna 64, 82
Fraser River, Harrison Rapids, salmon fry 459
Fresh-water mullet (*Agonostomus monticola*) 415–425
Description of larval forms 417
Occurrence and length 421, 424
Spawning 424
Frigate mackerel (*Auxis thazard*) at Point Judith, Rhode Island 336
Fundulus majalis, killifish, northern range 336

Page

Furunculosis in trout 555–564
drug sensitivity 556

Gehringer, Jack W.: OBSERVATIONS ON THE DEVELOPMENT OF THE ATLANTIC SAILFISH, *Istiophorus americanus* (CUVIER), WITH NOTES ON AN UNIDENTIFIED SPECIES OF ISTIOPHORID 139–171
Gempylus serpens, snake mackerel, food important to tuna 64, 67, 75, 79
Georges Bank, estimating groundfish population 265
Germo alalunga, skipjack 497
Glude, John B., and Harlan S. Spear: EFFECTS OF ENVIRONMENT AND HEREDITY ON GROWTH OF THE SOFT CLAM *Mya arenaria* 279–292
Gonyaulax monilata in Indian River, Fla 491
Graham, Herbert W., Clyde C. Taylor, and Henry B. Bigelow: CLIMATIC TRENDS AND THE DISTRIBUTION OF MARINE ANIMALS IN NEW ENGLAND 293–345
Great Lakes (Mich.), tagging trout 2
Green Bay lake herring (see lake herring) 88
economic importance of fishery 90
map of Green Bay 89
water level 90
Groundfish, Georges Bank, estimating abundance 265–278
Growth and development of the sailfish 150
Growth of marked Lake Michigan trout 36
in length 39–48
in weight 48–56
Growth of silver mullet 410
habitat of juvenile forms 413
Growth of soft clam 285
Growth rate of Lake herring 105, 107, 109
weight changes 104, 116
Gulf of Maine offshore water temperatures 302–307
temperature fluctuations in relation to fishes 293–345
Gulf of Mexico, new calanoid copepods of the families Aetideidae, Euchaetidae, and Stephidae 355–363
Gulf of Mexico, new genus and two new species of Tharibidae, with remarks on the status of the family 347–354
Gulf of Mexico, mortality of marine animals in association with *Gymnodinium brevis* 469
Gymnodinium brevis with unialgal cultures 472–496
red tide 469–496
Gymnura altavela, butterfly ray, northern range 335

Habitat and habits of larval forms of silver mullet 413
Heredity in soft shell clam, New England 279
Hida, Thomas S., and Joseph E. King: ZOOPLANKTON ABUNDANCE IN THE CENTRAL PACIFIC, PART II 365–395
Hippocampus hudsonius, sea horse, northern range 336
Hoven, Earl E., and Wilvan G. Van Campen: TUNAS AND TUNA FISHERIES OF THE WORLD: AN ANNOTATED BIBLIOGRAPHY, 1930–53 173–249

Page

Hypomesus pretiosus, silver smelt spawning, Pacific Coast 444

Ikehara, Isaac I., and Joseph E. King: COMPARATIVE STUDY OF FOOD OF BIGEYE AND YELLOWFIN TUNA IN THE CENTRAL PACIFIC 61–85
Istiophorids of the western north Atlantic 139
possible identity of specimens 169
list of specimens studied 170
sailfishes 139–171
Istiophorus orientalis 522–523
Istiompax marlina, black marlin 524

June, Fred C., and Heeny S. H. Yuen: YELLOWFIN TUNA SPAWNING IN THE CENTRAL EQUATORIAL PACIFIC 251–264
Juvenile sailfish 150

Karluk Lake and River 456–457
Key to Tharybidae family 349
Key to spearfishes of Central Pacific 499
King, Joseph E., and Isaac I. Ikehara: COMPARATIVE STUDY OF FOOD OF BIGEYE AND YELLOWFIN TUNA IN THE CENTRAL PACIFIC 61–85
King, Joseph E., and Thomas S. Hida: ZOOPLANKTON ABUNDANCE IN THE CENTRAL PACIFIC, PART II 365–395
King mackerel (*Scomberomorus regalis*) 336–337
Klamath River 452

Lake Michigan lake herring in Green Bay 87
Lake Michigan lake trout 1–59
Age determination from scales 9
Larvae of silver mullet in plankton tows 398
LARVAL FORMS OF THE FRESH-WATER MULLET (*Agonostomus monticola*) FROM THE OPEN OCEAN OFF THE BAHAMAS AND SOUTH ATLANTIC COAST OF THE UNITED STATES, by William W. Anderson 415–425
Length-weight relation in growth of lake trout 36
Leucichthys artedi (LeSueur), Lake Michigan herring 87
Leucichthys ascribed in Europe to *Argyrosomus* 87
LIFE HISTORY OF LAKE HERRING OF GREEN BAY, LAKE MICHIGAN, by Stanford H. Smith 87–138
Little tuna (*Euthynnus alletteratus*) in Barnstable 336
Lizard fish (*Synodus foetens*) 336
Lobster, Changes in abundance 325
fluctuations in landings 325
Loliginidae, food of tuna 66

MacGregor, John S.: FECUNDITY OF THE PACIFIC SARDINE (*Sardinops caerulea*) 427–449
Mackerel, fluctuations in landings 324
Makaira albida (Poey) 169
ample, blue marlin 532
audax, striped marlin 528
Mallotus catervarius, capelin, Pacific Coast 444
Marine animals, Changes in distribution 323
Marked Lake Michigan trout, release and recoveries 2
Marlin, blue 169
white 169
Meetinghouse Cove, New England, soft clam 287
Menhaden, Changes in abundance 333
Menticirrhus saxatilis, kingfish, northern range 337
METHOD OF ESTIMATING ABUNDANCE OF GROUNDFISH ON GEORGES BANK, A, by George A. Rounsefell 265–278
Michigan, Lake Trout, age determination 1–59
monticola, Agonostomus, fresh-water mullet 415
Mortality of marine animals in Gulf of Mexico 469
Mugil cephalus, experimental 486–492
striped-mullet, northern range 336–415
Mugil curema, silver mullet 397, 415
development of egg 399
growth 402, 410
occurrence 397, 410
spawning 410
Mugilidae, fresh-water mullet 415–425
Mullet, fresh-water 415
description 417
occurrence of larval and juvenile 424
spawning 424
Mya arenaria, soft clam 279–292

NEW CALANOID COPEPODS OF THE FAMILIES AETIDEIDAE, EUCHAETIDAE, AND STEPHIDAE FROM THE GULF OF MEXICO, by Abraham Fleminger 355–363
New England, soft shell clams 279–292
marine animals, distribution 293
NEW GENUS AND TWO NEW SPECIES OF THARYBIDAE (*Copepoda calanoida*) FROM THE GULF OF MEXICO, WITH REMARKS ON THE STATUS OF THE FAMILY, by Abraham Fleminger 347–354
Neothunnus macropterus (Temminck and Schlegel), distribution in Central Pacific 383
Neothunnus macropterus, food of yellowfin tuna 61, 79
North American Salmonidae 451–468
size and weight of eggs and fry 459
North Atlantic, Air and sea temperatures 294, 298, 301

OBSERVATIONS ON THE DEVELOPMENT OF THE ATLANTIC SAILFISH, *Istiophorus americanus* (CUVIER), WITH NOTES ON AN UNIDENTIFIED SPECIES OF ISTIOPHORID, by Jack W. Gehringer 139–171
OBSERVATIONS ON THE SPEARFISHES OF THE CENTRAL PACIFIC, by William F. Royce 497–554
Offatts Bayou (Galveston Bay), *Gonyaulax* cause of mortality 493
O-mark on Lake Michigan lake trout scales 26
Ommastrephidae and Loliginidae families, food for tuna fish 65, 75, 79
Oncorhynchus 451, 453
largest eggs among species 461
Oncorhynchus keta in Japanes waters 455
Otter trawlers from Boston on Georges Bank 266
amount of fishing in each depth zone 270

Page

Pacific, Central Equatorial, yellowfin tuna spawning 251–264
Pacific sardine 427–449
Pacific herring, *Clupea pallasi*, spawning 443
pacificus, Thaleichthys, eulachon spawning 443
Palmyra Island, zooplankton distribution 388
pallasi, Clupea 443
Paracalanina family 347
paraconcinna, Euchaeta, new species 358
Parundinella, new genus 348
Pink salmon, McClinton Creek 452, 453, 462
Pinnixa and *Polyonyx*, tube crabs 337
pretiosus, Hypomesus, silver smelt spawning, Pacific Coast 444
Prosopium, range in northwestern North America 87

Ray, Sammy M., and William B. Wilson: EFFECTS OF UNIALGAL AND BACTERIA-FREE CULTURES OF *Gymnodinium brevis* ON FISH, AND NOTES ON RELATED STUDIES WITH BACTERIA 469–496
Red tide 469, 492
Rounsefell, George A.: A METHOD OF ESTIMATING ABUNDANCE OF GROUNDFISH ON GEORGES BANK 265–278
Rounsefell, George A.: FECUNDITY OF NORTH AMERICAN SALMONIDAE 451–468
Royce, William F.: OBSERVATIONS ON THE SPEARFISHES OF THE CENTRAL PACIFIC 497–554

Sacramento River 452
Sagadahoc Bay, soft clams in 279
Georgetown Island 279
Sailfish, Atlantic (*Istiophorus americanus*) 139
food habits 164
growth 150
larvae 142–149
spawning 165
Sailfish, *Istiophorus orientalis* 139–171, 522
Salmo trutta, brown trout, experimental 555
fecundity of 451
treatment with sulfonamides and chloramphenicol 556–558
Salmonidae, fecundity of 451–468
Salvelinus fontinalis, brook trout, experimental 555
Salvelinus n. namaycush (Walbaum), lake trout 1–59
sarda sarda, bonito, northern range 336
Sardine, Fecundity of 427–449
Sardinops caerulea, Pacific sardine 427–449
age at first maturity 447
number of spawnings a year 440
relation between fecundity and age 439
Scale measurements of lake herring 92
Scales in age determination of Lake Michigan trout 1–59
Scomberomorus regalis, mackerel, northern range 336–337
Scolecithricella ctenopus (Giesbrecht) 347
Scripps Institution of Oceanography 365
Zooplankton studies, Central Pacific 365

Page

Sea bass (*Centropristes striatus*), Corea, Maine 337
Sea hare (*Tethys protea*), damage lobster trap, Woods Hole 337
Sea horse (*Hippocampus hudsonius*) Maine Coast 336
Sensitivity test with antibiotics 556,
with sulfonamides 558
Shortnose spearfish *Tetrapturus angustirostris* 520–522
Silver mullet, Atlantic Coast 397–414
development from larval to juvenile stage 402
growth 410
pigmentation 405
scales 407
spawning 410
Silver smelt (*Hypomesus pretiosus*, spawning 444
Size of yellowfin tuna at first spawning 257
Skeena River system 462
Smith, Stanford H.: LIFE HISTORY OF LAKE HERRING OF GREEN BAY, LAKE MICHIGAN 87–138
Snieszko, S. F., and G. L. Bullock: TREATMENT OF SULFONAMIDE-RESISTANT FURUNCULOSIS IN TROUT AND DETERMINATION OF DRUG SENSITIVITY 555–564
Sockeye salmon, Cultus Lake 453
Soft shell clam, New England Coast 279
South Atlantic Coast of U.S., fresh-water mullet 415–425
South Atlantic Coast, silver mullet 397–414
fishery investigations, sailfish 139–171
Spawning of fresh-water mullet (*Agonostomus monticola*) 424
surveys of habitat 415
Spawning of sailfish 165
Spawning of silver mullet 410
Spear, Harlan S., and John B. Glude: EFFECTS OF ENVIRONMENT AND HEREDITY ON GROWTH OF THE SOFT CLAM (*Mya arenaria*) 279–292
Spearfish, Tetrapturus angustirostris 520–522
Spearfishes of the Central Pacific 497–554
analysis of diagnostic characters 500
determination of allometric growth 502
key to spearfishes, Central Pacific 499
observations on species 518
Squid, food of yellowfin and bigeye tuna 67, 75, 79
Stephidae, Copepod from Gulf of Mexico 361
Stephos Scott 361
S. deichmannae, n. sp 363
Stizostediom vitreum, walleye scales compared with lake trout 6
Stomach contents, bigeye and yellowfin tuna 64
Striped bonito (*Euthynnus pelamis*), Barnstable Cape Cod 336
Striped marlin, *Makaira audax* 528
Striped mullet (*Mugil cephalus*), Green Bay, N.H. 336
Striped mummichog or killifish (*Fundulus majalis*,) Green Bay, N.H 336
Sulfonamide-resistant furunculosus in trout 555
Sulfonamides, Treatment of drug-resistant furunculosis 555
Swedish Deep-Sea Expedition Surveys in Pacific 365

Page

Swordfish, broadbill, *Xiphias gladius* 518
Synodus foetens, lizard fish, northern range 336

Tagged trout returns, Lake Michigan 2
Taylor, Clyde C., Henry B. Bigelow, and Herbert W. Graham: CLIMATIC TRENDS AND THE DISTRIBUTION OF MARINE ANIMALS IN NEW ENGLAND 293–345
Temperature changes in New England waters 294
Temperature in faunistic changes, New England 338
Temperatures over United States and Canada 294
Tethys protea, sea hare, found at Woods Hole 337
Tetrapturus angustirostris, shortnose spearfish 520–522
Tetrapturus belone Rafinisque 169
Thaleichthys pacificus, eulachon, spawning 443
Tharybidae, synoptic key 349
Two new species Tharybidae 347
Tharybis macrophthalma, northeastern Atlantic 347
Thunnus thynnus, tuna, northern range 336
Toxicity studies with bacteria 488–490
Trawlers, otter, Boston, for determining relative abundance of groundfish 266
fishing in each depth zone 270
TREATMENT OF SULFONAMIDE-RESISTANT FURUNCULOSIS IN TROUT AND DETERMINATION OF DRUG SENSITIVITY, BY S. F. Snieszko and G. L. Bullock 555–564
Trout, Lake Michigan 1–59
Tube crabs (*Pinnixa* and *Polyonyx*) rare in Woods Hole region 337
Tulane University collection, *Oregon* 170
Tuna abundance in Central Pacific, food of bigeye and yellowfin 61–85, 383
Tuna (*Thunnus thynnus*) in Gulf of Maine 336
TUNAS AND TUNA FISHERIES OF THE WORLD: AN ANNOTATED BIBLIOGRAPHY, 1930–53, by Wilvan Van Campen and Earl E. Hoven 173–249

Ulcer disease in trout, Experimental treatment 562
Undinella Sars 349
Unialgal cultures, Experiments with 469–495
Flavobacterium piscicida 489
Gymnodinium brevis 472–488
G. splendens 473
Prorocentrum sp 474, 475
U.S. National Museum collections of sailfish, taken by *Albatross, Fish Hawk, and Oregon* 170

VALIDITY OF AGE DETERMINATION FROM SCALES, AND GROWTH OF MARKED LAKE MICHIGAN LAKE TROUT, by Louella E. Cable 1–59
Van Campen, Wilvan G., and Earl E. Hoven: TUNAS AND TUNA FISHERIES OF THE WORLD: AN ANNOTATED BIBLIOGRAPHY 1930–53 173–249

Page

Venus mercenaria spawning temperature, New England 339
VESSELS:
Albatross, off New England 304, 307, 318
Alaska, Gulf of Mexico 139, 149, 170
Asterias, Woods Hole Oceanic Institute temperature readings in Massachusetts Bay 322
Carnegie Equatorial Pacific 365–395
Cavalieri 75
Charles H. Gilbert 75, 365
Fish Hawk 170
Grampus, New England 318
Halcyon 311
Hugh M. Smith 365, 381, 395
John R. Manning 75, 384, 497
Theodore N. Gill 139–140, 170, 397, 415, 417

Western Atlantic Istiophorids: Atlantic sailfish, white marlin, blue marlin, and spearfish 169
West coast of Florida fish mortality not caused by oxygen lack 493
White marlin 164, 169
Whiting, changes in abundance and distribution 329
Wilson, William B., and Sammy M. Ray: EFFECTS ON UNIALGAL AND BACTERIA-FREE CULTURES OF GYMNODINIUM BREVIS ON FISH 469–496
Wood borer (*Xylophaga dorsalis*), damage to lobster trap, Maine 337
Woods Hole, Mass., Mean water temperatures 345

Xiphias gladius, swordfish, broadbill 139, 164, 169, 518
Xylophaga dorsalis, wood borer, enemy of lobster traps, Maine 337

Yellowfin and bigeye tuna food 61
food varies with depth 67, 82
food varies with distance from land 69, 82
Yellowfin tuna, *Neothunnus macropterus*, food 383
YELLOWFIN TUNA SPAWNING IN THE CENTRAL EQUATORIAL PACIFIC, by Heeny S. H. Yuen and Fred C. June 251–264
Yellowtail flounder changes in abundance 334
Yuen, Heeny S. H., and Fred C. June: YELLOWFIN TUNA SPAWNING IN THE CENTRAL EQUATORIAL PACIFIC 251–264

ZOOPLANKTON ABUNDANCE IN THE CENTRAL PACIFIC, PART II, by Joseph E. King and Thomas S. Hida 365–395
distribution about an oceanic island 388
variations in system of currents 375, 377–382